(4					
		🛚 Career Institute)	(Reliable Nationa	al Entrance les	τ)
		ACAI	DEMIC SESSION : 2021-22		
		SAMPLE TE	ST PAPER (For XII to XIII	l Moving)	
ratio	n : 18 Min.			Max. Ma	arks : 96
me :		Арр	lication Form Number		
		Reg. Number : [2 0		
_		GE	NERAL INSTRUCTIONS)	
2. 3.	(9 to 16), M Blank paper gadgets in a Write your	lathematics(17 to 2 rs, clip boards, log ny form is not allow Name and Roll No.	24), each question carries 4 m tables, slide rule, calculators, wed.	ark. mobile or any other op of this booklet.	electronic
4. 5. 6. 7.	Before ans answer shee Do not forg the answer s No rough s the blank sp In case of a be final.	wering the paper, f et. et to mention your sheet. heets will be provide pace provided in the ny dispute, the answ	ill up the required details in t roll number neatly and clearly ded by the invigilators. All the question paper. wer filled in the OMR sheet a	the blank space provi y in the blank space p he rough work is to b available with the inst	ded in the rovided in be done in titute shall
4. 5. 6. 7.	Before ans answer shee Do not forg the answer s No rough s the blank sp In case of a be final.	wering the paper, f et. et to mention your sheet. heets will be provide pace provided in the ny dispute, the answ	ill up the required details in t roll number neatly and clearly ded by the invigilators. All the question paper. wer filled in the OMR sheet a	the blank space provi y in the blank space p he rough work is to b available with the inst	ded in the rovided in be done in titute shall
4. 5. 6. 7.	Before ans answer shee Do not forg the answer s No rough s the blank sp In case of a be final.	wering the paper, f et. et to mention your sheet. heets will be provide pace provided in the ny dispute, the answ	ill up the required details in t roll number neatly and clearly ded by the invigilators. All the question paper. wer filled in the OMR sheet a	the blank space provi y in the blank space p he rough work is to b available with the inst	ded in the rovided in be done in titute shall
4. 5. 6. 7.	Before ans answer shee Do not forg the answer s No rough s the blank sp In case of a be final.	wering the paper, f et. et to mention your sheet. heets will be provide pace provided in the ny dispute, the answ	ill up the required details in t roll number neatly and clearly ded by the invigilators. All the question paper. wer filled in the OMR sheet a MARKING CRITERIA	the blank space provi y in the blank space p he rough work is to b available with the inst Marks	ded in the rovided in be done in titute shall

PHYSICS

This section contains 8 multiple choice questions. Each question has four choices (1),(2),(3) and (4) out of which only one is correct

 A block A of mass 2 kg is hanging in a vertical plane with a spring of stiffness constant k = 100 N/m. A block B of mass 1 kg is kept on block A and the system is in equilibrium. Suddenly block B is removed. The amplitude of resulting SHM of A is

 (1) 5 cm
 (2) 10 cm
 (3) 15 cm



2. A particle is moving east-wards with a velocity of 4 m/s. In 10 seconds the velocity changes to 3 m/s northwards. The average acceleration in this time interval is

(1)
$$\frac{1}{2}$$
 m/s² towards north-east
(2) $\frac{1}{\sqrt{2}}$ m/s² towards north-west
(3) $\frac{1}{\sqrt{2}}$ m/s² towards north-east
(4) $\frac{1}{2}$ m/s² towards north-west

3. A block of mass 1 kg just remains in equilibrium with the vertical wall of a cart accelerating uniformly with 20 m/s² as shown. The co-efficient of friction between block and wall is ($g = 10 \text{ m/s}^2$)

$$(1) 0.1 (2) 0.2 (3) 0.5 (4) 1$$

4. Two balls of mass M = 9 g and m = 3 g are attached by massless threads AO and OB. The length AB is 1 m. They are set in rotational motion in a horizontal plane about a vertical axis at O with constant angular velocity ω . The ratio of length

AO and
$$OB\left(\frac{AO}{OB}\right)$$
 for which the tension in threads are same

will be

(1) $\frac{1}{3}$ (2) 3 (3) $\frac{2}{3}$ (4) $\frac{3}{2}$





R-NET (2020)



- 5. A and B are two concentric metallic hollow spheres. If A is given a charge q while B is earthed as shown in figure, then
 (1) charge density of A and B are same
 - (2) field inside and outside A is zero
 - (3) field between A and B is not zero
 - (4) field inside and outside B is zero
- 6. Each cell has emf ε and internal resistance *r* in the figure. Find the current through resistance *R*

(1)
$$\frac{4\varepsilon}{r}$$
 (2) $\frac{3\varepsilon}{r}$

(3)
$$\frac{\varepsilon}{r}$$
 (4) zero

7. Light is incident normally on face AB of a prism as shown in figure. A liquid of refractive index μ is placed on face AC of the prism. The prism is made of glass of refractive index 3/2. The limits of μ for which total internal reflection cannot takes place on face AC is

(1)
$$\frac{3\sqrt{3}}{4} > \mu > \frac{\sqrt{3}}{2}$$
 (2) $\mu > \frac{3\sqrt{3}}{4}$ (3) $\mu > \sqrt{3}$ (4) $\mu < \frac{\sqrt{3}}{2}$

8. A uniform but time varying magnetic field is present in a circular region of radius R. The magnetic field is perpendicular and into the plane of the paper and the magnitude of the field is increasing at a constant rate α . There is a straight conducing rod of length 2R placed as shown in the figure. The magnitude of induced emf across the rod is

(1)
$$\pi R^2 \alpha$$
 (2) $\frac{\pi R^2 \alpha}{2}$ (3) $\frac{R^2 \alpha}{\sqrt{2}}$ (4) $\frac{\pi R^2 \alpha}{4}$

SAMPLE TEST PAPER (FOR XII TO XIII MOVING)











CHEMISTRY

This section contains 8 multiple choice questions. Each question has four choices (1),(2),(3) and (4) out of which only one is correct

- **9.** A 4.8 molal aqueous solution of ethyl alcohol is supplied, what is the mole fraction of ethyl alcohol in the solution.
 - (1) 0.079 (2) 0.014 (3) 0.067 (4) 0.09
- 10. Which of the following metal on treatment with NaOH (aq.) will liberate H_2 gas.
 - (1) Zn (2) Mg (3) K (4) Cu
- A solution prepared at 25°C by mixing 10 mL of 1 molar HCl and 10 mL of 2 molar NaOH has a pH of
 - (1) 14.0 (2) 0.3 (3) 13 (4) 13.7
- 12. In which of the following, Ea for backward reaction is greater than Ea forward reaction
 - (i) $A \xrightarrow{Ea=50 \text{ Kcal}} B$; $\Delta H = -10 \text{ kcal}$
 - (ii) $A \xrightarrow{Ea=50 \text{ Kcal}} B$; $\Delta H = +10 \text{ kcal}$
 - (iii) A + 10 K cal \rightarrow B; Ea = 50 kcal
 - (iv) $A 10 \text{ K cal} \rightarrow B$; Ea = 50 kcal

Which is correct–

- (1) (i), (ii), (iii) (2) (i), (iv) (3) (ii), (iii), (iv) (4) (i), (iii)
- 13. First compound for Xe synthesized was-
 - (1) $[Xe F^+][Xe PtF_5]^-$ (2) $[Xe O_2]$ (3) $Xe [PtF_6]$ (4) $O_2[Xe F_6]$

R-NET (2020)



14. In the reaction shown below, the major product(s) formed is/are



15. The major product of the following reaction is :



16. The major product obtained in the following reaction is :-





MATHEMATICS

This section contains 8 multiple choice questions. Each question has four choices (1),(2),(3) and (4) out of which only one is correct

17. If $f(x) = x \cdot \frac{(a^{1/x} - a^{-1/x})}{(a^{1/x} + a^{-1/x})}, x \neq 0 (a > 1), f(0) = 0$ then

- (1) f is differentiable at x = 0
- (2) f is not differentiable at x = 0
- (3) f is not continuous at x = 0
- (4) None of these
- 18. Let z be a complex number such that $5z + 3\overline{z} = 8 + 2i$, then arg(z) is-

(1)
$$2n\pi + \frac{3\pi}{4}; n \in I$$

(2) $2n\pi + \frac{\pi}{2}; n \in I$
(3) $2n\pi + \frac{\pi}{4}; n \in I$
(4) $2n\pi + \frac{5\pi}{4}; n \in I$

19. Let $\vec{a} = \hat{i} - \hat{k}$, $\vec{b} = x\hat{i} + \hat{j} + (1-x)\hat{k}$ and $\vec{c} = y\hat{i} + x\hat{j} + (1+x-y)\hat{k}$. Then $[\vec{a}\vec{b}\vec{c}]$ depends on:

- (1) only y
 (2) only x
 (3) both x and y
 (4) Neither x nor y
- **20.** Let 3x y 8 = 0 be the equation of tangent to a parabola at the point (7, 13). If the focus of the parabola is at (-1, -1), its directrix is
 - (1) x 8y + 19 = 0(2) 8x + y + 19 = 0(3) 8x - y + 19 = 0(4) x + 8y + 19 = 0
- 21. If r_1 and r_2 are the radii of smallest and largest circles which passes through (5, 6) and touches the circle $(x 2)^2 + y^2 = 4$, then r_1r_2 is
 - (1) $\frac{4}{41}$ (2) $\frac{41}{4}$ (3) $\frac{5}{41}$ (4) $\frac{41}{6}$

R-NET (2020)

6/8



- 22. The tangent and normal drawn to the curve $y = x^2 x + 4$ at P(1, 4) cut the x axis at A and B respectively. If the length of the subtangent drawn to the curve at P is equal to the length of the subnormal then the area of the triangle PAB in square unit is :
 - (1) 4 (2) 8 (3) 16 (4) 32

23.
$$\int \frac{\sec x (1 + \tan x) dx}{(e^{-x} + \sec x)} = f(x) + C, \text{ where } f(0) = \ln 2, \text{ then } f\left(\frac{\pi}{4}\right) \text{ is } -$$

(1)
$$\ell n \left(1 + e^{\frac{\pi}{4}} \sqrt{2} \right)$$
 (2) $\ell n (\sqrt{2})$ (3) $\ell n \left(2\sqrt{2} \right)$ (4) $\ell n \left(\frac{e^{\frac{\pi}{4}}}{\sqrt{2}} + 1 \right)$

- **24.** The last two digits of the number 9^{200} are :
 - (1) 81 (2) 43 (3) 29 (4) 01



Space for rough work



R-NET (SAMPLE PAPER) (XII to XIII moving students)

ANSWER KEY																				
Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	2	4	3	1	3	4	2	4	1	1	4	2	3	1	4	2	2	3	4	4
Que.	21	22	23	24																
Ans.	2	3	1	4																